

<https://helda.helsinki.fi>

COVID-19 and SIC (!)

Nevzorov, Ilja

2020-09

Nevzorov , I , Tulamo , R , Albäck , A & Lassila , R 2020 , ' COVID-19 and SIC (!) ' , Journal of Vascular Surgery , vol. 72 , no. 3 , pp. 1148-1150 . <https://doi.org/10.1016/j.jvs.2020.05.047>

<http://hdl.handle.net/10138/333342>

<https://doi.org/10.1016/j.jvs.2020.05.047>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Journal Pre-proof

COVID-19 and SIC (!)

Ilya Nevzorov, PhD, Riikka Tulamo, MD, PhD, Anders Albäck, MD, PhD, Riitta Lassila, MD, PhD



PII: S0741-5214(20)31303-3

DOI: <https://doi.org/10.1016/j.jvs.2020.05.047>

Reference: YMVA 11368

To appear in: *Journal of Vascular Surgery*

Received Date: 14 April 2020

Accepted Date: 22 May 2020

Please cite this article as: Nevzorov I, Tulamo R, Albäck A, Lassila R, COVID-19 and SIC (!), *Journal of Vascular Surgery* (2020), doi: <https://doi.org/10.1016/j.jvs.2020.05.047>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2020 Published by Elsevier Inc. on behalf of the Society for Vascular Surgery.

COVID-19 and SIC (!)

Ilja Nevzorov, PhD¹, Riikka Tulamo, MD, PhD², Anders Albäck, MD, PhD² and Riitta Lassila, MD, PhD^{3,*}.

¹ Faculty of Medicine, University of Helsinki, Helsinki, Finland.

² Department of Vascular Surgery, Helsinki University Hospital, University of Helsinki, Helsinki, Finland.

³ Coagulation Disorders Unit, University of Helsinki, Departments of Haematology and Clinical Chemistry (HUSLAB Laboratory Services), Comprehensive Cancer Center, Helsinki University Hospital and Research Program in Systems Oncology, Faculty of Medicine, University of Helsinki, Helsinki, Finland; Helsinki University, Faculty of Medicine, Research Program in Oncology, Helsinki, Finland; Aplagon Oy, Helsinki, Finland.

* Corresponding Author: riitta.lassila@hus.fi

To the Editor,

Accurate risk stratification tools are paramount for optimal disease management. Patients with cardiovascular conditions, diabetes and cancer are most susceptible to COVID-19 complications leading to poor outcome¹. These systemic diseases relate to enhanced fibrin formation and thromboinflammation. Indeed, severity of peripheral occlusive arterial disease correlates with the levels of both fibrinogen and its turnover measure D-dimer². In severe COVID-19-infection, elevation of D-dimer and sepsis-induced coagulopathy (SIC), predicts poor prognosis. The incidence of venous thromboembolism (VTE) in patients with severe COVID-19 pneumonia is 25% (!)³. Furthermore, endothelial injury inherent to vascular procedures may predispose to coagulopathy in COVID-19. The benefit of low-molecular-weight heparin therapy is protection of critically ill patients against VTE, as well as its putative anti-inflammatory properties. Pulmonary

embolism (PE), triggered by severe infection, may be masked by the symptoms and signs of hypoxia in COVID-19. We advocate these considerations for vascular specialists.

A large retrospective Chinese cohort study¹ demonstrated that the fibrin turnover-measure D-dimer exceeding 1 µg/mL on admission was associated with an increased risk of in-hospital death (OR 20, 95% CI 6.5-61.56, $P < .0001$) in COVID-19 patients. Another retrospective study⁴ assessed the benefits of anticoagulation on 28-day mortality, which does not appear to differ between heparin users (22%) and nonusers (mortality rates 30.3% vs 29.7%, respectively). However, patients with 6-fold D-dimer levels (3 µg/mL) to normal clearly benefited from anticoagulation, translating to lower mortality (32.8% vs 52.4%, $P = .017$). Therefore, D-dimer levels on admission are particularly useful for risk stratification in COVID-19 patients (Fig. 1).

Another important predictor of mortality is the International Society of Thrombosis and Haemostasis (ISTH) SIC-score⁵, which includes prothrombin time (ratio >1.5), platelet count ($<100 \times 10^9/l$) and sequential organ function assessment (SOFA-score). In the above-mentioned study³, patients with ISTH SIC-score of ≥ 4 treated with anticoagulation showed again lower 28-day mortality rates than the untreated ones (40% vs 64%; $P=.029$).

To guarantee the best outcomes for patients we suggest that all medical professionals, including vascular specialists, adhere to ISTH guidelines on recognition and management of coagulopathy in COVID-19 based on D-dimer and SIC-scores as major prognostic factors⁶ (Fig. 1).

References:

1. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054-62.
2. Lassila R, Peltonen S, Lepäntalo M, Saarinen O, Kauhanen P, Manninen V. Severity of peripheral atherosclerosis is associated with fibrinogen and degradation of cross-linked fibrin. *Arterioscler Thromb Vasc Biol*. 1993;13:1738-1742.
3. Cui S, Chen S, Li X, Liu S, Wang F. Prevalence of venous thromboembolism in patients with severe novel coronavirus pneumonia. *J Thromb Haemost*. 2020;00:1-4. Epub 2020 Apr 9.
4. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost*. 2020 May;18(5):1094-1099. Epub 2020 March 27.
5. Iba T, Nisio MD, Levy JH, Kitamura N, Thachil J. New criteria for sepsis-induced coagulopathy (SIC) following the revised sepsis definition: a retrospective analysis of a nationwide survey. *BMJ Open*. 2017;7(9):e017046.
6. Thachil J, Tang N, Gando S, Falanga A, Cattaneo M, Levi M, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. 2020 May;18(5):1023-1026. Epub 2020 March 25.

Figure 1. Outline of the algorithm for the management of coagulopathy in COVID-19 based on D-dimer and SIC-score. DVT – deep vein thrombosis, PE – pulmonary embolism, LMWH – low molecular weight heparin, SIC – sepsis-induced coagulopathy.

